

Weak Bruhat order on the set of faces of the permutahedra.

Patricia PALACIOS¹ and María O. RONCO²

Abstract. *Given a Coxeter system (W, S) , we define a partial order on the set of right coclasses $\mathcal{P}^{(W, S)} := \{W_J \circ w : \text{for } W_J \text{ a parabolic subgroup of } W\}$, which extends the weak Bruhat order of the group W . When W is the group of permutations of n elements, we get a partial order on the set of faces of the permutahedra, which induces a partial order on the set of planar rooted trees with $n + 1$ leaves. We use these orders to describe associative algebra structures on the vector spaces spanned by the faces of permutahedra on one side, and on the vector space spanned by the planar rooted trees, on the other one. The associative products of both algebras may be described as the sum of three different operations, also given by the weak Bruhat order, which give to our examples the structure of dendriform trialgebras.*

Introduction. Let S_n be the group of permutations of n elements. In [MR], C. Malvenuto and C. Reutenauer defined a Hopf algebra structure on the vector space $k[S_\infty] := \bigoplus_{n \geq 0} k[S_n]$, where k is a field and $k[S_n]$ is the group algebra of the symmetric group. The associative product of $k[S_\infty]$ is defined using the natural action of the Solomon algebra Sol_∞ (cf. [GKLLR] and [LR1]) on $k[S_\infty]$. This algebra is widely studied in [DHT].

From another point of view, J.-L. Loday introduced in [L2] the notion of dendriform algebra. In the same work he described free dendriform algebras in terms of planar binary rooted trees. As a particular case, he showed that the vector space $k[\mathcal{Y}_\infty]$, spanned by all planar binary rooted trees, has a natural structure of free dendriform algebra generated by one element. For a fixed field k , there exists a canonical functor from the category of dendriform algebras into the category of associative algebras (over k). The associative product of $k[\mathcal{Y}_\infty]$ is also described in [L2] and [HNT].

In [LR1] and [LR2], the authors used the existence of a canonical surjection from S_n into the set $\mathcal{Y}_n$ of planar binary trees with $n + 1$ leaves, and a definition of the

¹Depto. de Matemáticas, FCEN, Univ. de Buenos Aires. Pab. I, Ciudad Universitaria (1428) Buenos Aires, Argentina. E-mail : ppalacio@bigua.dm.uba.ar

²Depto. de Matemática, CBC, Univ. de Buenos Aires. Pab. III Ciudad Universitaria (1428) Buenos Aires, Argentina. E-mail : mronco@mate.dm.uba.ar

canonical generators of the Solomon algebra in terms of the weak Bruhat order of S_n , to prove that :

1. The associative structure of the Malvenuto-Reutenauer algebra comes from a dendriform algebra structure on $k[S_\infty]$, which may be defined in terms of two elementary associative operations $/$ and $\backslash$ and the weak Bruhat order.
2. The weak Bruhat order of S_n induces a partial order on $\mathcal{Y}_n$, in such a way that the associative structure of the free dendriform algebra $k[\mathcal{Y}_\infty]$ can also be described in terms of this order, and are induced by the dendriform algebra structure of $k[S_\infty]$.

In more recent works (cf. [LR3] and [LR4]), the vector space $k[\mathcal{T}_\infty]$ spanned by the set of all planar rooted trees is described as the free object on one element for a new type of algebras, called dendriform trialgebras.

The purpose of this paper is to:

1. Extend the weak Bruhat order of a Coxeter group W to the set $\mathcal{P}^{(W,S)} := \{W_J \circ w : \text{for } W_J \text{ a parabolic subgroup of } W\}$.
2. Prove that this order defines a structure of dendriform trialgebra on the vector space $k[\mathcal{P}_\infty]$ spanned by the all the posets $\mathcal{P}_n$ associated to the symmetric groups.
3. Show that there exists a natural surjection from the set $\mathcal{P}_n$ into the set $\mathcal{T}_n$ of planar rooted trees with $n + 1$ leaves, such that the weak Bruhat order of $\mathcal{P}_n$ induces a partial order on $\mathcal{T}_n$. This order defines a structure of free dendriform trialgebra on the vector space $k[\mathcal{T}_\infty]$, generated by all planar rooted trees, which coincides with the one defined in [LR3].

The first point is described in Section 1.

In Section 2, we give a description of the poset $\mathcal{P}_n$ in terms of surjective maps from $\{1, \dots, n\}$ into $\{1, \dots, r\}$, for positive integers n and r . This description permit us to construct subsets $SH(n_0, \dots, n_r)$ of $\mathcal{P}_n$, which generalize the well-known notion of shuffles in S_n . We study the orders $\subseteq$ and $\leq_B$ in this new context, define a structure of dendriform trialgebra on $k[\mathcal{P}_\infty]$ and show that this structure may be obtained in terms of the generalized weak Bruhat order.

Section 3 is devoted to see that the orders on $\mathcal{P}_n$ induce orders on the set $\mathcal{T}_n$ of planar rooted trees with $n + 1$ leaves and to describe the free dendriform trialgebra structure of $k[\mathcal{T}_\infty]$ in terms of the weak Bruhat order defined in this context.

1 Weak Bruhat order

In this Section we define an order on the set of right coclasses $\mathcal{P}^{(W,S)} := \{W_J \circ w : \text{for } W_J \text{ a parabolic of } W\}$, which differs from the canonical inclusion order and extends the weak Bruhat order of the group W . For the elementary definitions and results about Coxeter groups, Solomon algebra and weak Bruhat order we refer to [Bo], [Hu], [S] and [BBHT]. We shall deal only with finite Coxeter systems.

- Notation 1**
1. *Given a Coxeter system (W, S) , we denote by $l(w)$ the length of an element $w \in W$. The element ω_S is the unique element of maximal length in W .*
 2. *Given a subset $J \subseteq S$, we denote by W_J the standard parabolic subgroup of W generated by J . The pair (W_J, J) is also a Coxeter system.*
 3. *For a subset J of S , X_J denotes the Solomon subset of elements which have no descent at J . We recall that $X_J := \{w \in W \mid l(w \circ s) > l(w), \forall s \in J\}$, where $\circ$ denotes the product of the group W .*

Let us recall the definition of the weak Bruhat order on W .

Definition 2 *Let (W, S) be a Coxeter system, the weak Bruhat order on W is defined by $x \leq_B x'$ if $x' = y \circ x$, with $l(x') = l(y) + l(x)$.*

The group W equipped with the weak Bruhat order $\leq_B$ is a partially ordered set. The minimal element of $(W, \leq_B)$ is the identity element 1_W of the group, and its maximal element is ω_S .

Given a subset $J \subseteq S$, there exist unique elements $\xi_J \in X_J$ and $\omega_J \in W_J$ such that $\omega_S = \xi_J \circ \omega_J$. It is easy to check that ω_J is the maximal element of (W_J, J) , and that ξ_J is the longest element of X_J .

The following result is proved in [LR2].

Lemma 3 *Let (W, S) be a Coxeter system and let $J \subseteq S$.*

1. *The subset X_J of W verifies that $X_J = \{w \in W : w \leq_B \xi_J\}$.*
2. *The subgroup W_J of W is the set of elements which are smaller or equal than ω_J .*

Definition 4 *Let (W, S) be a Coxeter system. The Coxeter poset of (W, S) is the set $P^{(W, S)}$ of right coclasses modulo the parabolic standard subgroups, ordered by the inclusion $\subseteq$.*

The following result follows immediately from [Bo] Ch. IV, p. 37, ex. 3.

Lemma 5 *The set $P^{(W, S)}$ may be described as $P^{(W, S)} := \{W_J \circ w \text{ with } J \subseteq S \text{ and } w \in X_J^{-1}\}$.*

For $J = \emptyset$ one has that $X_\emptyset = W$. So, we may think the group W as embedded in $P^{(W, S)}$, via the map $w \mapsto W_\emptyset \circ w$.

The family of subsets $\mathcal{P}_r^{(W, S)} := \{W_J \circ w : |J| = r\}$, for $0 \leq r \leq |S|$, defines a graduation on the set $\mathcal{P}^{(W, S)}$.

Let us define an order, different from $\subseteq$, on the set $P^{(W, S)}$.

In [BBHT] the authors proved that for any subset $J \subseteq S$ and $s_0 \in S \setminus J$, it holds that $X_{J \cup \{s_0\}} \circ X_J^{W_{J \cup \{s_0\}}} = X_J$, where $X_J^{W_{J \cup \{s_0\}}}$ is the set of elements in $W_{J \cup \{s_0\}}$ which have no descent at J .

Using this equality we get the following definition

Definition 6 *Let $J \subseteq S$. For any $s \in S \setminus J$, the result above implies that there exists an element $\alpha_{J, s} \in X_J^{W_{J \cup \{s\}}} = X_J \cap W_{J \cup \{s\}}$ such that $\xi_J = \xi_{J \cup \{s\}} \circ \alpha_{J, s}$.*

Definition 7 (Weak Bruhat order on $P^{(W, S)}$) *Let (W, S) be a Coxeter system. The weak Bruhat order on the set $P^{(W, S)}$ is the transitive relation generated by the conditions:*

1. *For $J \subseteq S$, $s \in S \setminus J$ and $w \in X_{J \cup \{s\}}^{-1}$, it holds that $W_J \circ w <_B W_{J \cup \{s\}} \circ w$.*
2. *For $J \subseteq S$, $s \in S \setminus J$ and $w \in X_{J \cup \{s\}}^{-1}$, it holds that $W_{J \cup \{s\}} \circ w <_B W_J \circ (\alpha_{J, s}^{-1} \circ w)$.*

To check that this gives a partial order relation it suffices to see that, given a coclass $W_J \circ w$ it is impossible to get, applying several times the relations 1 and 2, a sequence of coclasses that ends at $W_J \circ w$. Note that applying 2 we get $W_J \circ w <_B W_K \circ z$ only for $w <_B z$. So, we may restrict our proof to a sequence of coclasses obtained by applying only the first relation. But, applying only 1, we get that $W_J \circ w <_B W_K \circ z$ implies J is strictly contained in K .

The weak Bruhat order of W induces an order on the subset $\{W_\emptyset \circ w \text{ with } w \in W\} \subseteq P^{(W,S)}$. The proof of the following Lemma is straightforward.

Lemma 8 *Let $w <_B z$ be two elements of W . The coclass $W_\emptyset \circ w$ is smaller than $W_\emptyset \circ z$ for the weak Bruhat order.*

Let us point out that the result above is false if we replace $\emptyset$ by any $J \subseteq S$. As an example, consider the symmetric group S_4 with the set of generators $\{s_1, s_2, s_3\}$, where s_i is the permutation that exchanges i and $i + 1$. The coclasses $W_{\{s_2, s_3\}} \circ 1_{S_4}$ and $W_{\{s_2, s_3\}} \circ s_1$ are not comparable for the weak Bruhat order.

Lemma 9 *Let $K \subseteq S$. If z and w are two elements of W such that $z \in X_K^{-1}$ and $\omega_K \circ z \leq_B w$, then $W_K \circ z \leq_B W_J \circ w$, for any J with $w \in X_J^{-1}$.*

Proof. We prove the lemma by induction on the number of elements of K .

Suppose $|K| = 0$. We have that $z \leq_B \omega_K \circ z \leq_B w$, and Lemma 8 shows that $W_\emptyset \circ z \leq_B W_\emptyset \circ w$. Now, by applying several times the relation 1 of Definition 7, we get that $W_\emptyset \circ w \leq_B W_J \circ w$, and the result holds.

For $|K| \geq 1$, we have that $K = K' \cup \{s\}$ with $|K'| = |K| - 1$. The relation 2 of Definition 7 states that $W_K \circ z <_B W_{K'} \circ (\alpha_{K',s}^{-1} \circ z)$.

From the equalities $\omega_S = \xi_{K'} \circ \omega_{K'} = \xi_K \circ \omega_K$, and $\xi_{K'} = \xi_K \circ \alpha_{K',s}$, we get $\omega_K = \alpha_{K',s} \circ \omega_{K'}$. Since it holds that $\omega_{K'} \circ \alpha_{K',s}^{-1} = \omega_{K'}^{-1} \circ \alpha_{K',s}^{-1} = \omega_K^{-1} = \omega_K$, we get

$$\omega_{K'} \circ (\alpha_{K',s}^{-1} \circ z) = \omega_K \circ z < w.$$

Applying a recursive argument, $W_K \circ z <_B W_{K'} \circ (\alpha_{K',s}^{-1} \circ z) <_B W_J \circ w$. $\diamond$

The following Theorem describes the relationship between the orders $\subseteq$ and $\leq$ on $P^{(W,S)}$.

Theorem 10 *Let (W, S) be a Coxeter system. For any $J \subseteq S$ and any $w \in X_J^{-1}$, the sets $\{W_K \circ z \in P^{(W,S)} : W_K \circ z \subseteq W_J \circ w\}$ and $\{W_K \circ z \in P^{(W,S)} : W_\emptyset \circ w \leq_B W_K \circ z \leq_B W_\emptyset \circ (\omega_J \circ w)\}$ are equal.*

Proof. Observe that $W_K \circ z \subseteq W_J \circ w$ if, and only if $K \subseteq J$ and $w \leq_B z \leq_B \omega_J \circ w$. Since $w \leq_B z$, we have that $W_\emptyset \circ w \leq_B W_\emptyset \circ z \leq_B W_K \circ z$. Now, $W_K \circ z \subseteq W_J \circ w$ implies that $\omega_K \circ z \leq_B \omega_J \circ w$. Applying Lemma 9, we get that $W_K \circ z \leq_B W_\emptyset \circ (\omega_J \circ w) = \omega_J \circ w$.

We have proved that the set $\{W_K \circ z \in P^{(W,S)} : W_K \circ z \subseteq W_J \circ w\}$ is contained in $\{W_K \circ z \in P^{(W,S)} : W_\emptyset \circ w \leq_B W_K \circ z \leq_B W_\emptyset \circ (\omega_J \circ w)\}$.

Suppose that $W_\emptyset \circ w \leq_B W_K \circ z \leq_B W_\emptyset \circ (\omega_J \circ w)$. From the first inequality it is immediate that $w \leq_B z$, from the second one we get that $z \leq_B \omega_J \circ w$. There exists $y_0 \in W_J$ such that $z = y_0 \circ w$. To check that $W_K \circ z \subseteq W_J \circ w$ we only need to prove that $K \subseteq J$. But for any $x \in W_K$, it holds that $W_\emptyset \circ w \leq_B W_K \circ (x \circ z) \leq_B W_\emptyset \circ (\omega_J \circ w)$, which implies that $w \leq_B x \circ z \leq_B \omega_J \circ w$. So, $x \circ z = y_x \circ w$, for some $y_x \in W_J$; consequently $x = y_x \circ y_0^{-1}$. We get that $W_K \subseteq W_J$, which implies (cf. [Bo]) that $K \subseteq J$. $\diamond$

2 Symmetric groups

Let $(S_n, \circ)$ be the group of permutations of n elements. The group S_n is a Coxeter group generated by $n - 1$ transpositions $s_1, \dots, s_{n-1}$, where s_i is the permutation that exchanges i and $i + 1$. We shall denote the set $\{s_1, \dots, s_{n-1}\}$ by $\mathcal{S}_n$.

The longest element of S_n is the permutation $\omega_n := s_1 \circ s_2 \circ s_1 \circ \dots \circ s_{n-2} \circ \dots \circ s_1 \circ s_n \circ s_{n-1} \circ \dots \circ s_1$.

2.1 The posets associated to the symmetric group

Let $(n_1, \dots, n_r)$ be partition of n . Given elements $\sigma_i \in S_{n_i}$, for $1 \leq i \leq r$, the permutation $\sigma_1 \times \dots \times \sigma_r$ in S_n is defined as:

$$\sigma_1 \times \dots \times \sigma_r := (\sigma_1(1), \dots, \sigma_1(n_1), \sigma_2(1) + n_1, \dots, \sigma_r(n_r) + n_1 + \dots + n_{r-1}).$$

We denote by $S_{n_1, \dots, n_r}$ the subgroup of S_n which is the image of the embedding $S_{n_1} \times \dots \times S_{n_r} \hookrightarrow S_n$.

Let $(n_1, \dots, n_r)$ be a partition of n , and let J be the set $\mathcal{S}_n \setminus \{s_{n_1}, s_{n_1+n_2}, \dots, s_{n_1+\dots+n_{r-1}}\}$. It is immediate to check that the standard parabolic subgroup W_J is the subgroup $S_{n_1, \dots, n_r}$ of S_n . Moreover, the longest element of $S_{n_1, \dots, n_r}$ is $\omega_{n_1, \dots, n_r} := \omega_{n_1} \times \dots \times \omega_{n_r}$.

Given a sequence of non-negative integers $n_1, \dots, n_r$ such that $\sum_{i=1}^r n_i = n$, a $(n_1, \dots, n_r)$ -shuffle is a permutation $\sigma \in S_n$ verifying that:

$$\sigma(1) < \dots < \sigma(n_1), \dots, \sigma(n_1 + \dots + n_{r-1} + 1) < \dots < \sigma(n).$$

The set of all $(n_1, \dots, n_r)$ -shuffles in S_n is denoted by $Sh(n_1, \dots, n_r)$.

Remark 11 1. Let σ be an element of S_n and s_i a transposition in S_n . It is easily verified that $\text{length}(s_i \circ \sigma) = \text{length}(\sigma) + 1$ if, and only if, $\sigma^{-1}(i) < \sigma^{-1}(i+1)$.

2. Let $(n_1, \dots, n_r)$ be a partition of n . The assertion above implies that the set $Sh(n_1, \dots, n_r)$ coincides with the set X_J described in the first section, for

$$J = S_n \setminus \{s_{n_1}, s_{n_1+n_2}, \dots, s_{n_1+\dots+n_{r-1}}\}.$$

The longest element of $Sh(n_1, \dots, n_r)$ (cf. [LR2]) is the permutation $\xi_{n_1, \dots, n_r}$ defined as:

$$\xi_{n_1, \dots, n_r}(k) := k - \sum_{i=1}^{j-1} n_i + \sum_{i=j+1}^r n_i, \text{ for } \sum_{i=1}^{j-1} n_i < k \leq \sum_{i=1}^j n_i, \text{ for } 1 \leq j \leq r.$$

Notation 12 In order to simplify notation, we denote by $\mathcal{P}_n$ the set $P^{(S_n, S_n)}$.

The description of the set $P^{(W, S)}$ given in Section 1 and the results above imply that for any right coclass $W_J \circ \tau \in \mathcal{P}_n$, there exist a unique partition $(n_1, \dots, n_r)$ of n and a unique element $\sigma \in Sh(n_1, \dots, n_r)^{-1}$ such that $W_J \circ \tau = S_{n_1, \dots, n_r} \circ \sigma$.

For any partition $(n_1, \dots, n_r)$ of n and any $1 \leq k \leq r-1$, we denote by $\alpha_{n_1, \dots, n_r}^k$ the permutation such that $\xi_{n_1, \dots, n_r} = \xi_{n_1, \dots, n_k+n_{k+1}, \dots, n_r} \circ \alpha_{n_1, \dots, n_r}^k$.

Recall that we identify an element σ of S_n with the coclass $S_{1, \dots, 1} \circ \sigma$. So we have an embedding of partially ordered sets $(S_n, \leq_B) \hookrightarrow (\mathcal{P}_n, \leq_B)$.

Let us give another description of the sets $\mathcal{P}_n$, which makes it easier to deal with.

Lemma 13 There exists a bijection between the set $\mathcal{P}_n$ and the set of all surjective maps $\gamma : \{1, \dots, n\} \rightarrow \{1, \dots, r\}$, with $r \geq 1$.

Proof. Let $x = S_{n_1, \dots, n_r} \circ \sigma$ be an element of $\mathcal{P}_n$.

Consider the map $\Psi(x) : \{1, \dots, n\} \rightarrow \{1, \dots, r\}$ defined by:

$$\Psi(x)(i) := j, \text{ if } n_1 + \dots + n_{j-1} < \sigma(i) \leq n_1 + \dots + n_j.$$

Observe that the map Ψ is bijective, the inverse map is defined by

$\Psi^{-1}(\gamma) = S_{n_1, \dots, n_r} \circ \sigma$, where $Im(\gamma) = \{1, \dots, r\}$, n_i is the number of elements in $\gamma^{-1}(i)$ and $\sigma(j) := n_1 + \dots + n_{\gamma(j)-1} + k$, for $\gamma^{-1}(\gamma(j)) = \{i_1 < \dots < i_{n_j}\}$ and $i_k = j$. $\diamond$

Note that the image of the element $S_{1,1,\dots,1} \circ \sigma$ under Ψ is the permutation σ .

So, $\mathcal{P}_n = \{\gamma : \{1, \dots, n\} \rightarrow \{1, \dots, r\} \text{ surjective, for some } r \geq 1\}$.

Fix the set $\mathcal{P}_0 := \{(0)\}$. Let $\mathcal{P}_\infty$ denotes the disjoint union of all the sets $\mathcal{P}_n$, for $n \geq 0$. The usual composition of maps defines a multiplication on $\mathcal{P}_\infty$:

$$\gamma \circ \delta := \begin{cases} \gamma \circ \delta & \text{if } \gamma \in \mathcal{P}_r \text{ and } Im(\delta) = \{1, \dots, r\} \\ (0) & \text{otherwise.} \end{cases}$$

From now on, we shall denote an element $\gamma \in \mathcal{P}_n$ by its image $\gamma = (\gamma(1), \dots, \gamma(n))$. If γ and δ are permutations, then their composite coincides with their product in the symmetric group S_n .

Remark 14 *Note that any element $\gamma \in \mathcal{P}_n$, with $n \geq 1$, may be written as $\gamma = \gamma' \circ \sigma$ with $\sigma \in S_n$ and γ' a non-decreasing map. The element γ' is unique, while there exist many permutations σ verifying the equality. However there exists a unique σ of minimal length, which is precisely the permutation defined in Lemma 13 verifying $\Psi^{-1}(\gamma) = S_{n_1, \dots, n_r} \circ \sigma$.*

For $1 \leq r \leq n$, the subset $\mathcal{P}_{n,r} := \{\gamma \in \mathcal{P}_n : Im(\gamma) = \{1, \dots, r\}\}$ corresponds to the subset $\mathcal{P}_r^{(S_n, S_n)}$ defined in Section 1.

There exists an embedding $\mathcal{P}_n \times \mathcal{P}_m \hookrightarrow \mathcal{P}_{n+m}$, given by:

$$(\gamma, \delta) \mapsto (\gamma(1), \dots, \gamma(n), \delta(1) + r, \dots, \delta(m) + r),$$

where $\gamma \in \mathcal{P}_{n,r}$. The image of (γ, δ) under this embedding is denoted by $\gamma \times \delta$. Note that $\gamma \times (0) = \gamma = (0) \times \gamma$, for all $\gamma \in \mathcal{P}_\infty$.

For $n \geq 1$, let us denote by t_i the element of $\mathcal{P}_{n,n-1}$ defined by:

$$t_i(j) := \begin{cases} j & \text{for } 1 \leq j \leq i \\ j-1 & \text{for } i < j \leq n. \end{cases}$$

Remark 15 The product of the element t_j with the transpositions s_i , which span the symmetric group, verify the following relationships:

$$s_i \circ t_j = \begin{cases} t_j \circ s_i & \text{if } i < j - 1 \\ t_{j-1} \circ s_j \circ s_{j-1} & \text{if } i = j - 1 \\ t_{j+1} \circ s_j \circ s_{j+1} & \text{if } i = j \\ t_j \circ s_{i+1} & \text{if } i > j. \end{cases}$$

2.2 Shuffles and wedges on $\mathcal{P}_n$

We proceed to define a notion of shuffle in $\mathcal{P}_n$ which extends the definition of shuffle in S_n .

Definition 16 Let $\gamma \in \mathcal{P}_n$ and $(n_1, \dots, n_r)$ be a partition of n , we say that γ is a $(n_1, \dots, n_r)$ -shuffle in $\mathcal{P}_n$ if

$$\gamma(1) < \dots < \gamma(n_1), \quad \dots, \quad \gamma(n_1 + \dots + n_{r-1} + 1) < \dots < \gamma(n).$$

We denote by $SH(n_1, \dots, n_r)$ the subset of $(n_1, \dots, n_r)$ -shuffles in $\mathcal{P}_n$. The following result is a generalized version of [Bo], Ch. IV, p. 37, ex. 3, to the set $\mathcal{P}_n$.

Lemma 17 For any element $\gamma \in \mathcal{P}_n$, and any $0 \leq p \leq n$, there exist unique elements $\gamma_1 \in \bigcup_{\substack{1 \leq i \leq p \\ 1 \leq j \leq n-p}} SH(i, j)$, $\delta_1 \in \mathcal{P}_{p,i}$ and $\delta_2 \in \mathcal{P}_{n-p,j}$ such that $\gamma = \gamma_1 \circ (\delta_1 \times \delta_2)$.

Proof. Let γ be an element of $\mathcal{P}_{n,r}$, there exist unique integers $1 \leq i \leq p$ and $1 \leq j \leq n - p$ and unique bijective order-preserving morphisms

$$\phi_1 : \gamma(\{1, \dots, p\}) \rightarrow \{1, \dots, i\} \text{ and } \phi_2 : \gamma(\{p+1, \dots, n\}) \rightarrow \{1, \dots, j\}.$$

Define $\delta_1 : (\phi_1(\gamma(1)), \dots, \phi_1(\gamma(p)))$, $\delta_2 := (\phi_2(\gamma(p+1)), \dots, \phi_2(\gamma(n)))$, and

$\gamma_1 := (\phi_1^{-1}(1), \dots, \phi_1^{-1}(i), \phi_2^{-1}(1), \dots, \phi_2^{-1}(j))$. It is easy to check that $\gamma_1 \in SH(i, j)$ and that the elements are unique. $\diamond$

Definition 18 Let $\gamma_0, \dots, \gamma_k$ be a family of elements of $\mathcal{P}_\infty$, with $\gamma_i \in \mathcal{P}_{n_i, r_i}$ or $\gamma_i = (0)$, and let ω be an element of $SH(r_0, \dots, r_k)$. The wedge of $\gamma_0, \dots, \gamma_k$ over ω is the element of $\mathcal{P}_{n+k}$ given by:

$$\bigvee_{\omega} (\gamma_0, \dots, \gamma_k) := (\omega \circ (\gamma_0 \times \dots \times \gamma_k) \times 1_1) \circ z(n_0, \dots, n_k),$$

where $z(n_0, \dots, n_k) := (1, \dots, n_0, n_0+1, n_0+1, \dots, n_0+n_1, n_0+1, \dots, n_0+1, n_0+\dots+n_{k-1}+1, \dots, n)$ and $n := \sum_{i=0}^k n_i$.

For instance, observe that $1^n := (1, 1, \dots, 1) : \{1, \dots, m\} \rightarrow \{1\}$ is the wedge $\bigvee_{(0)}((0), \dots, (0))$.

Proposition 19 *Given $\gamma \in \mathcal{P}_{n,r}$ there exist unique elements γ_i in $\mathcal{P}_{n_i, r_i}$, with $k + \sum_{i=0}^k n_i = n$ and $r - 1 = \sum_{i=0}^k r_i$, and $\omega \in SH(r_0, \dots, r_k)$ such that $\gamma = \bigvee_{\omega}(\gamma_0, \dots, \gamma_k)$.*

Proof. Let $\gamma^{-1}(r) = \{j_1 < \dots < j_k\}$. Define the integers n_i , for $0 \leq i \leq k$, as follows:

$$n_0 := j_1 - 1, \quad \dots, \quad n_l := j_{l+1} - j_l - 1, \quad \dots, \quad n_k := n - j_k.$$

It is easy to check that $\gamma = (\alpha \times 1_1) \circ z(n_0, \dots, n_k)$, with $\alpha \in \mathcal{P}_{n-k, r-1}$. From Lemma 13, we get that there exist unique $\gamma_0 \in \mathcal{P}_{n_0}, \dots, \gamma_k \in \mathcal{P}_{n_k}$ and $\omega \in SH(r_0, \dots, r_k)$ such that $\alpha = \omega \circ (\gamma_0 \times \dots \times \gamma_k)$.

The argument above proves the existence of the decomposition. It is clear that α and $n_0, \dots, n_k$ are unique. Lemma 13 implies the unicity of the γ_i 's and ω . $\diamond$

Definition 20 *Let $\omega \in SH(r_0, \dots, r_k)$ and let $1 \leq j \leq \sum_{i=0}^k r_i$. We associate to ω and j a family of positive integers $(s_0^j, \dots, s_k^j)$, with $s_i^j = r_i$ or $s_i^j = r_i - 1$, and a shuffle $\omega_j \in SH(s_0^j, \dots, s_k^j)$ in the following way:*

1. *If $t_j \circ \omega \in SH(r_0, \dots, r_k)$, then $(s_0^j, \dots, s_k^j) := (r_0, \dots, r_k)$ and $\omega_j := t_j \circ \omega$.*
2. *If $t_j \circ \omega \notin SH(r_0, \dots, r_k)$, then*

$$s_i^j := \begin{cases} r_i & \text{if } \{j, j+1\} \not\subseteq \omega(\{R_{i-1} + 1, \dots, R_i\}), \\ r_i - 1 & \text{if } \{j, j+1\} \subseteq \omega(\{R_{i-1} + 1, \dots, R_i\}), \end{cases}$$

where $R_l := r_0 + \dots + r_l$. The element ω_j is the $(s_0^j, \dots, s_k^j)$ -shuffle in $\mathcal{P}_{\sum s_i^j}$ such that, for $0 \leq i \leq k$ and $1 \leq l \leq s_i^j$, $\omega_j(s_0^j + \dots + s_{i-1}^j + l) :=$

$$\begin{cases} \omega(R_{i-1} + l) & \text{if } s_i^j = r_i, \\ \omega(R_{i-1} + l) & \text{if } s_i^j = r_i - 1 \text{ and } \omega(R_{i-1} + l) \leq j, \\ \omega(R_{i-1} + l + 1) - 1 & \text{if } s_i^j = r_i - 1 \text{ and } \omega(R_{i-1} + l) > j. \end{cases}$$

Lemma 21 *Let $\omega \in SH(r_0, \dots, r_k)$ and $\gamma_i \in \mathcal{P}_{n_i, r_i}$, for $0 \leq i \leq k$.*

For $1 \leq j \leq r := \sum_{i=0}^k r_i + 1$, it holds that:

1. If $j < r-1$ and $t_j \circ \omega \in SH(r_0, \dots, r_k)$, then $t_j \circ V_\omega(\gamma_0, \dots, \gamma_k) = V_{t_j \circ \omega}(\gamma_0, \dots, \gamma_k)$.
2. If $j < r-1$ and $t_j \circ \omega \notin SH(r_0, \dots, r_k)$, then $t_j \circ V_\omega(\gamma_0, \dots, \gamma_k) = V_{\omega_j}(\gamma_0^j, \dots, \gamma_k^j)$,
 where $\gamma_i^j := \begin{cases} \gamma_i & \text{if } s_i^j = r_i, \\ t_{j_i} \circ \gamma_i & \text{if } s_i^j = r_i - 1, \end{cases}$
 for $\omega^{-1}(j) = r_0 + \dots + r_{i-1} + j_i$, with $0 \leq i \leq k$.

Proof. The proof follows straightforward, using the definition above. $\diamond$

2.3 Inclusion order and Weak Bruhat order on $\mathcal{P}_n$

Translating the definition of inclusion order and weak Bruhat order to the set of surjective maps $\gamma : \{1, \dots, n\} \rightarrow \{1, \dots, r\}$, we get the following result.

Lemma 22 1. Let $\gamma \in \mathcal{P}_{n,r}$ and $\delta \in \mathcal{P}_{n,s}$. It holds that $\gamma \subseteq \delta$ in $\mathcal{P}_n$ if, and only if, there exists a non-decreasing map $\rho \in \mathcal{P}_{r,s}$ such that $\delta = \rho \circ \gamma$.

2. The weak Bruhat order $\leq_B$ on the set $\mathcal{P}_n$ is the transitive relation spanned by the conditions:

- (a) If $\gamma^{-1}(i) < \gamma^{-1}(i+1)$ for some $1 \leq i \leq r$ then $\gamma <_B t_i \circ \gamma$,
- (b) If $\gamma^{-1}(i) > \gamma^{-1}(i+1)$ for some $1 \leq i \leq r$ then $t_i \circ \gamma <_B \gamma$,

for $\gamma \in \mathcal{P}_{n,r}$, where $\gamma^{-1}(i) < \gamma^{-1}(i+1)$ means that any element j of $\gamma^{-1}(i)$ verifies $j < k$ for all $k \in \gamma^{-1}(i+1)$.

Proof. The proof of the first point is immediate.

For the second one, it is easily seen that the weak Bruhat order $\leq_B$ defined on the elements of $\mathcal{P}_n$, seen as right coclasses, is the transitive relation spanned by the following conditions:

1. $S_{n_1, \dots, n_r} \circ \sigma <_B S_{n_1, \dots, n_k + n_{k+1}, \dots, n_r} \circ \sigma$.
2. $S_{n_1, \dots, n_k + n_{k+1}, \dots, n_r} \circ \sigma <_B S_{n_1, \dots, n_r} \circ (\alpha_{n_1, \dots, n_r}^k)^{-1} \circ \sigma$.

for any partition $(n_1, \dots, n_r)$ of n , any $1 \leq k \leq r-1$ and any $\sigma \in Sh(n_1, \dots, n_k + n_{k+1}, \dots, n_r)$. Translating these conditions from the elements of type $S_{n_1, \dots, n_r} \circ \sigma$ to the surjective maps $\{1, \dots, n\} \rightarrow \{1, \dots, r\}$ the assertion follows. $\diamond$

Remark 23 1. Given $\gamma \in \mathcal{P}_n$ it holds that $1_n \leq_B \gamma \leq_B \omega_n$, where 1_n is the identity element of S_n and $\omega_n = (n, n-1, \dots, 1)$.

2. Suppose $\gamma = \tau \circ \sigma$, with τ a non-decreasing map and $\sigma \in S_n$. If

$\gamma \in SH(n_1, \dots, n_r)$, then $\sigma \in Sh(n_1, \dots, n_r)$.

3. Suppose $\gamma \leq_B \gamma'$ in $\mathcal{P}_n$. If $\gamma'(j) < \gamma'(k)$ for some $1 \leq j < k \leq n$, then $\gamma(j) < \gamma(k)$. Conversely, if $\gamma(j) > \gamma(k)$ for some $1 \leq j < k \leq n$, then $\gamma'(j) > \gamma'(k)$.

Proposition 24 It holds that

$$SH(n_1, \dots, n_r) = \{\omega \in \mathcal{P}_n \text{ such that } \omega \leq_B \xi_{n_1, \dots, n_r}\}.$$

Proof. Suppose $\omega \in \mathcal{P}_n$ is such that $\omega \leq_B \xi_{n_1, \dots, n_r}$. The Remark above implies that $\omega \in SH(n_1, \dots, n_r)$.

Conversely, suppose $\omega \in SH(n_1, \dots, n_r)$. We have that $\omega = \tau \circ \sigma$, with τ a non-decreasing map and $\sigma \in Sh(n_1, \dots, n_r)$, which may be supposed of minimal length. But τ may be written as a product $\tau = t_{j_k} \circ \dots \circ t_{j_1}$, with $j_1 \leq j_2 \leq \dots \leq j_k$, for unique integers $j_1, \dots, j_k$. If $k = 0$, the result follows from the Remark above.

For $k = 1$, we have two possibilities:

1. If $\sigma^{-1}(j_1 + 1) < \sigma^{-1}(j_1)$, then $\omega <_B \sigma \leq_B \xi_{n_1, \dots, n_r}$.
2. If $\sigma^{-1}(j_1) < \sigma^{-1}(j_1 + 1)$, then $\omega <_B s_{j_1} \circ \sigma$. Since $\omega \in SH(n_1, \dots, n_r)$, it holds that there exists $0 \leq l \leq r-1$ such that $\sigma^{-1}(j_1) \leq n_1 + \dots + n_l < \sigma^{-1}(j_1 + 1)$, which implies that $s_{j_1} \circ \sigma \in Sh(n_1, \dots, n_r)$. So, $\omega < s_{j_1} \circ \sigma \leq \xi_{n_1, \dots, n_r}$.

For $k > 1$, let $\gamma := t_{j_{k-1}} \circ \dots \circ t_{j_1} \circ \sigma$. The element $s_{j_k} \circ \gamma$ verifies that $\omega <_B s_{j_k} \circ \gamma$, $s_{j_k} \circ \gamma \in SH(n_1, \dots, n_r)$ and, using Remark 15, we get that

$$s_{j_k} \circ \gamma = t_{l_{k-1}} \circ \dots \circ t_{l_1} \circ \sigma', \quad \text{for some } \sigma' \in S_n.$$

So, applying recursive hypothesis to $s_{j_k} \circ \gamma$, we get that $\omega <_B s_{j_k} \circ \gamma \leq_B \xi_{n_1, \dots, n_r}$. $\diamond$

Proposition 25 Let $\gamma_0, \dots, \gamma_k$ and $\delta_0, \dots, \delta_k$ be families of elements in $\mathcal{P}_\infty$ verifying that $\gamma_i \leq_B \delta_i$ in $\mathcal{P}_{n_i}$. It holds that $\gamma_0 \times \dots \times \gamma_k \leq_B \delta_0 \times \dots \times \delta_k$.

Proof. Clearly, it suffices to show that

$$\gamma_0 \times \cdots \times \gamma_k \leq_B \gamma_0 \times \cdots \gamma_{i-1} \times \delta_i \times \gamma_{i+1} \times \cdots \times \gamma_k,$$

for all $0 \leq i \leq k$.

Fix $0 \leq i \leq k$, and suppose $\gamma_l \in \mathcal{P}_{n_l, r_l}$, for $0 \leq l \leq k$. The proof may be restricted to the following two cases:

1. When $\gamma_i^{-1}(j) < \gamma_i^{-1}(j+1)$, for some $1 \leq j \leq r_i$, and $\delta_i = t_j \circ \gamma_i$.
2. When $\delta_i^{-1}(j) > \delta_i^{-1}(j+1)$, for some j , and $\gamma_i = t_j \circ \delta_i$.

The proofs of both cases are analogous, so we give the proof of the first one.

If $\gamma_i^{-1}(j) < \gamma_i^{-1}(j+1)$, for some $1 \leq j \leq r_i$ and $\delta_i = t_j \circ \gamma_i$, then

$$(\gamma_0 \times \cdots \times \gamma_k)^{-1}(r_0 + \cdots + r_{i-1} + j) < (\gamma_0 \times \cdots \times \gamma_k)^{-1}(R_{i-1} + j + 1), \quad \text{and}$$

$$\gamma_0 \times \cdots \times \gamma_{i-1} \times \delta_i \times \gamma_{i+1} \times \cdots \times \gamma_k = t_{R_{i-1}+j} \circ (\gamma_0 \times \cdots \times \gamma_k),$$

where $R_{i-1} := r_0 + \cdots + r_{i-1}$.

So, $\gamma_0 \times \cdots \times \gamma_k <_B \gamma_0 \times \cdots \times \gamma_{i-1} \times \delta_i \times \gamma_{i+1} \times \cdots \times \gamma_k$. $\diamond$

Lemma 26 *Let $\gamma_0, \dots, \gamma_k$ be a family of elements in $\mathcal{P}_\infty$ such that $\gamma_i \in \mathcal{P}_{n_i, r_i}$, and let ω and ω' be elements of $SH(r_0, \dots, r_k)$ such that $\omega \leq_B \omega'$. It holds that:*

$$\omega \circ (\gamma_0 \times \cdots \times \gamma_k) \leq_B \omega' \circ (\gamma_0 \times \cdots \times \gamma_k).$$

Proof. Again, it suffices to consider the cases:

1. $\omega^{-1}(j) < \omega^{-1}(j+1)$ for some j , and $\omega' = t_j \circ \omega$.
2. $(\omega')^{-1}(j) > (\omega')^{-1}(j+1)$ for some j , and $\omega = t_j \circ \omega'$.

Since the proofs of the cases are very similar, we give the proof of the second one.

Suppose $(\omega')^{-1}(j) > (\omega')^{-1}(j+1)$ for some j , and $\omega = t_j \circ \omega'$. Since $\omega \in SH(r_0, \dots, r_k)$, there must exist $0 \leq l \leq k-1$ such that $(\omega')^{-1}(j+1) \subseteq \{1, \dots, n_0 + \cdots + n_l\}$ and $(\omega')^{-1}(j) \subseteq \{n_0 + \cdots + n_l + 1, \dots, n_0 + \cdots + n_k\}$. So, we get that

$$(\omega' \circ (\gamma_0 \times \cdots \times \gamma_k))^{-1}(j) > (\omega' \circ (\gamma_0 \times \cdots \times \gamma_k))^{-1}(j+1),$$

and $\omega \circ (\gamma_0 \times \cdots \times \gamma_k) = t_j \circ \omega' \circ (\gamma_0 \times \cdots \times \gamma_k)$. The argument above implies that:

$$\omega \circ (\gamma_0 \times \cdots \times \gamma_k) \leq_B \omega' \circ (\gamma_0 \times \cdots \times \gamma_k). \quad \diamond$$

Corollary 27 *Let $\gamma_0, \dots, \gamma_k$ be elements in $\mathcal{P}_\infty$ with $\gamma_i \in \mathcal{P}_{n_i, r_i}$. If $\omega \leq_B \omega'$ in $SH(r_0, \dots, r_k)$, then $\bigvee_\omega(\gamma_0, \dots, \gamma_k) \leq_B \bigvee_{\omega'}(\gamma_0, \dots, \gamma_k)$.*

Lemma 28 *Let $\gamma_i \in \mathcal{P}_{n_i, r_i}$, for $0 \leq i \leq k$, and let $\omega \in SH(r_0, \dots, r_k) \cap S_{r-1} = Sh(r_0, \dots, r_k)$.*

1. *If $\gamma_l \leq_B t_j \circ \gamma_l$ and $\omega(r_0 + \dots + r_{l-1} + j + 1) = \omega(r_0 + \dots + r_{l-1} + j) + 1$, for some $0 \leq l \leq k$ and $1 \leq j \leq r_l$, then:*

$$\bigvee_\omega(\gamma_0, \dots, \gamma_k) \leq_B \bigvee_{\omega'}(\gamma_0, \dots, t_j \circ \gamma_l, \dots, \gamma_k),$$

where $\omega' := t_{\omega(r_0 + \dots + r_{l-1} + j)} \circ \omega$.

2. *If $t_j \circ \gamma_l \leq_B \gamma_l$ and $\omega(r_0 + \dots + r_{l-1} + j + 1) = \omega(r_0 + \dots + r_{l-1} + j) + 1$, for some $0 \leq l \leq k$ and $1 \leq j \leq r_l$, then:*

$$\bigvee_{\omega'}(\gamma_0, \dots, t_j \circ \gamma_l, \dots, \gamma_k) \leq_B \bigvee_\omega(\gamma_0, \dots, \gamma_k),$$

where $\omega' := t_{\omega(r_0 + \dots + r_{l-1} + j)} \circ \omega$.

Proof. The proof is immediate, using that $t_{\omega(r_0 + \dots + r_{l-1} + j)} \circ \omega \in SH(r_0, \dots, r_k)$. $\diamond$

2.4 Dendriform trialgebra structure on the vector space spanned by $\mathcal{P}_\infty$

We want to extend some well-known results for the shuffles in S_n to our generalized shuffles in $\mathcal{P}_n$. Let us begin with the associativity of the shuffle.

Proposition 29 (Associativity of SH) *Given non-negative integers n, m, r the following equality holds:*

$$SH(n, m, r) = \bigcup_{1 \leq j \leq n+m} SH(j, r) \circ (SH(n, m) \times 1_r) = \bigcup_{1 \leq k \leq m+r} SH(n, k) \circ (1_n \times SH(m, r))$$

Proof. From Lemma 13, we get that for any $\gamma \in SH(n, m, r)$ there exist unique integers $1 \leq j \leq n + m$ and $1 \leq k \leq m + r$, and unique elements $\gamma_1 \in SH(j, r)$, $\gamma_2 \in SH(n, k)$, $\delta_1 \in \mathcal{P}_{n+m}$, $\delta_2 \in \mathcal{P}_r$, $\tau_1 \in \mathcal{P}_n$ and $\tau_2 \in \mathcal{P}_{m+r}$, such that

$$\gamma = \gamma_1 \circ (\delta_1 \times \delta_2) = \gamma_2 \circ (\tau_1 \times \tau_2).$$

To end the proof note that:

(i) Since $\gamma \in SH(n, m, r)$ and $\gamma_1 \in SH(j, r)$, the element δ_1 must belong to $SH(n, m)$ and δ_2 must be equal to 1_r .

(ii) The fact that $\gamma \in SH(n, m, r)$ and that $\gamma_2 \in SH(n, k)$ imply that $\tau_1 = 1_n$ and $\tau_2 \in SH(m, r)$. $\diamond$

For $n, m \geq 0$, the set $SH(n, m)$ is the disjoint union of the following three subsets:

$$\begin{aligned} SH^{\succ}(n, m) &:= \{\gamma \in SH(n, m) : \gamma(n) < \gamma(n+m)\}, \\ SH^{\bullet}(n, m) &:= \{\gamma \in SH(n, m) : \gamma(n) = \gamma(n+m)\}, \\ SH^{\prec}(n, m) &:= \{\gamma \in SH(n, m) : \gamma(n) > \gamma(n+m)\}. \end{aligned}$$

Lemma 30 *For $n, m \geq 0$, the sets $SH^{\succ}(n, m)$, $SH^{\bullet}(n, m)$ and $SH^{\prec}(n, m)$ may be described in terms of the weak Bruhat order as follows:*

1. $SH^{\succ}(n, m) = \{\gamma \in \mathcal{P}_{n+m} : \gamma \leq_B \xi_{n, m-1} \times 1_1\},$
2. $SH^{\bullet}(n, m) = \{\gamma \in \mathcal{P}_{n+m} : z(n-1, m-1, 0) \leq_B \gamma \leq_B (\xi_{n-1, m-1} \times 1_1) \circ z(n-1, m-1, 0)\},$
3. $SH^{\prec}(n, m) = \{\gamma \in \mathcal{P}_{n+m} : z(n-1, m) \leq_B \gamma \leq_B \xi_{n, m}\},$

where $z(n-1, m-1, 0) = (1, \dots, n-1, n+m-1, n, \dots, n+m-1)$ and $z(n-1, m) = (1, \dots, n-1, n+m, n, \dots, n+m-1)$.

Proof. The proof of the formulas follows from Proposition 24. We give the proof of the first equality, the others may be checked in a similar way.

If $\gamma \in SH^{\succ}(n, m)$, then $\gamma = \gamma_1 \times 1_1$, with $\gamma_1 \in SH(n, m-1)$. By Proposition 24, we get that $\gamma_1 \leq_B \xi_{n, m-1}$. It implies that $\gamma \leq_B \xi_{n, m-1} \times 1_1$.

Conversely, let $\gamma \in \mathcal{P}_{n+m}$ be such that $\gamma \leq_B \xi_{n, m-1} \times 1_1$. It is easily seen that $\gamma = \gamma_1 \times 1_1$, with $\gamma_1 \leq_B \xi_{n, m-1}$. So, $\gamma \in SH^{\succ}(n, m)$. $\diamond$

Remark 31 *It is not difficult to check that the following identities hold:*

1. $\bigcup_{i=1}^{m+r} SH^{\succ}(n, i) \circ (1_n \times SH^{\succ}(m, r)) = \bigcup_{j=1}^{n+m} SH^{\succ}(j, r) \circ (SH(n, m) \times 1_r),$
2. $\bigcup_{i=1}^{m+r} SH^{\prec}(n, i) \circ (1_n \times SH^{\prec}(m, r)) = \bigcup_{j=1}^{n+m} SH^{\prec}(j, r) \circ (SH^{\succ}(n, m) \times 1_r),$
3. $\bigcup_{i=1}^{m+r} SH^{\prec}(n, i) \circ (1_n \times SH(m, r)) = \bigcup_{j=1}^{n+m} SH^{\prec}(j, r) \circ (SH^{\prec}(n, m) \times 1_r),$

4. $\bigcup_{i=1}^{m+r} SH^\bullet(n, i) \circ (1_n \times SH^\bullet(m, r)) = \bigcup_{j=1}^{n+m} SH^\bullet(j, r) \circ (SH^\bullet(n, m) \times 1_r),$
5. $\bigcup_{i=1}^{m+r} SH^\succ(n, i) \circ (1_n \times SH^\bullet(m, r)) = \bigcup_{j=1}^{n+m} SH^\bullet(j, r) \circ (SH^\succ(n, m) \times 1_r),$
6. $\bigcup_{i=1}^{m+r} SH^\bullet(n, i) \circ (1_n \times SH^\succ(m, r)) = \bigcup_{j=1}^{n+m} SH^\bullet(j, r) \circ (SH^\prec(n, m) \times 1_r),$
7. $\bigcup_{i=1}^{m+r} SH^\bullet(n, i) \circ (1_n \times SH^\prec(m, r)) = \bigcup_{j=1}^{n+m} SH^\prec(j, r) \circ (SH^\bullet(n, m) \times 1_r).$

The following definition is given in [LR3], it generalizes the definition of dendriform algebra introduced by J.-L. Loday in [L2].

Definition 32 *Let k be a field. A dendriform trialgebra over k is a vector space V equipped with three k -linear maps $\succ, \cdot, \prec : V \otimes_k V \rightarrow V$ verifying the following identities:*

1. $v \succ (w \succ u) = (v * w) \succ u,$
2. $v \succ (w \prec u) = (v \succ w) \prec u,$
3. $v \prec (w * u) = (v \prec w) \prec u,$
4. $v \cdot (w \cdot u) = (v \cdot w) \cdot u,$
5. $v \succ (w \cdot u) = (v \succ w) \cdot u,$
6. $v \cdot (w \succ u) = (v \prec w) \cdot u,$
7. $v \cdot (w \prec u) = (v \cdot w) \prec u,$

for all v, w and u in V , where $v * w := v \succ w + v \cdot w + v \prec w$.

If $(V, \succ, \cdot, \prec)$ is a tridendriform algebra, then $(V, *)$ is an associative algebra over k . Let $k[\mathcal{P}_\infty]$ be the free k vector space spanned by the set $\mathcal{P}_\infty$. We define operations $\succ, \cdot$ and $\prec$ on $k[\mathcal{P}_\infty]$ as follows:

1. $\gamma \succ \delta := \sum_{\omega \in SH^\succ(r, s)} \omega \circ (\gamma \times \delta),$
2. $\gamma \cdot \delta := \sum_{\omega \in SH^\bullet(r, s)} \omega \circ (\gamma \times \delta),$
3. $\gamma \prec \delta := \sum_{\omega \in SH^\prec(r, s)} \omega \circ (\gamma \times \delta),$

for $\gamma \in \mathcal{P}_{n,r}$ and $\delta \in \mathcal{P}_{m,s}$.

Remark 31 implies that $k[\mathcal{P}_\infty]$ is a dendriform trialgebra over k . Moreover, it is easy to see that the associative product $*$ is given by the formula:

$$\gamma * \delta = \sum_{\omega \in SH(r,s)} \omega \circ (\gamma \times \delta), \quad \text{for } \gamma \in \mathcal{P}_{n,r} \quad \text{and} \quad \delta \in \mathcal{P}_{m,s}.$$

Theorem 33 *The operations of the dendriform trialgebra $k[\mathcal{P}_\infty]$ may be described in terms of the weak Bruhat order as follows:*

$$\begin{aligned} \gamma \succ \delta &= \sum_{\gamma \times \delta \leq_B \omega \leq_B (\xi_{r,s-1} \times 1_1) \circ (\gamma \times \delta)} \omega, \\ \gamma \cdot \delta &= \sum_{z(r-1,s-1,0) \circ (\gamma \times \delta) \leq_B \omega \leq_B (\xi_{r-1,s-1} \times 1_1) \circ z(r-1,s-1,0) \circ (\gamma \times \delta)} \omega, \\ \gamma \prec \delta &= \sum_{z(r-1,s) \circ (\gamma \times \delta) \leq_B \omega \leq_B \xi_{r,s}(\gamma \times \delta)} \omega, \\ \gamma * \delta &= \sum_{\gamma \times \delta \leq_B \omega \leq_B \xi_{r,s} \circ (\gamma \times \delta)} \omega, \end{aligned}$$

where $\gamma \in \mathcal{P}_{n,r}$, $\delta \in \mathcal{P}_{m,s}$.

Proof. It suffices to apply Lemmas 26 and 30. ◇

3 Planar rooted trees

In this paper, a *tree* will always be a finite planar non-empty oriented connected graph t without loops, and such that for any vertex of t there are at least two incoming edges and exactly one outgoing edge.

For $n \geq 0$, let $\mathcal{T}_n$ denote the set of planar rooted trees with $n + 1$ leaves:

$$\mathcal{T}_0 = \{\downarrow\}, \mathcal{T}_1 = \left\{ \begin{array}{c} \swarrow \quad \nwarrow \\ \bullet \\ \downarrow \end{array} \right\}, \mathcal{T}_2 = \left\{ \begin{array}{c} \swarrow \quad \nwarrow \\ \bullet \\ \swarrow \quad \nwarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \end{array}, \begin{array}{c} \swarrow \quad \nwarrow \\ \bullet \\ \downarrow \quad \nwarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \end{array}, \begin{array}{c} \swarrow \quad \nwarrow \\ \bullet \\ \swarrow \quad \nwarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \end{array} \right\}.$$

The *degree* of a tree t is n when t belongs to $\mathcal{T}_n$. We denote the degree of t by $|t|$. A tree t is called *binary* if each vertex of t has exactly two incoming edges. We denote by $\mathcal{Y}_n$ the subset of binary trees of $\mathcal{T}_n$. Note that the elements of $\mathcal{Y}_n$ are the trees which have exactly $n + 1$ leaves and n vertices.

The tree ς_n is the element of $\mathcal{T}_n$ which has exactly one vertex.

The *wedge* of planar rooted trees $t^0 \in \mathcal{T}_{n_0}, \dots, t^r \in \mathcal{T}_{n_r}$, is the tree $\bigvee(t^0, \dots, t^r)$ of degree $\sum_{i=0}^r n_i + r - 1$, obtained by joining the roots of $t^0, \dots, t^r$ to a new vertex and creating a new root. For any tree $t \neq \downarrow$, there exist unique trees $t^0, \dots, t^r$ such that $t = \bigvee(t^0, \dots, t^r)$.


We have that $\mathcal{T}_n = \bigsqcup_{n_0 + \dots + n_r + r - 1 = n} \mathcal{T}_{n_1} \times \dots \times \mathcal{T}_{n_r}$, for $n \geq 1$.

We denote by $\mathcal{T}_\infty$ the graded set of all planar trees $\bigcup_{n \geq 0} \mathcal{T}_n$.

3.1 Orders on $\mathcal{T}_n$

We are going to define two orders $\subseteq$ and $\leq_B$ on the set of planar trees $\mathcal{T}_n$. The first one is the opposite of the order described by V. Ginzburg and M. Kapranov in [GK]. In the next section we show that both orders are induced by the orders $\subseteq$ and $\leq_B$ on the set $\mathcal{P}_n$.

Let t be a tree and let e be an internal edge of t . We denote by t_e the tree obtained from t by contracting e into a point. For instance, if $t =$



Definition 34 Let $\subseteq$ be the order on $\mathcal{T}_n$ transitively generated by the relation:

$$t \subseteq t_e, \text{ for any internal edge } e \text{ of } t.$$

Note that the binary trees of $\mathcal{T}_n$ are the minimal elements for $\subseteq$, while the element ς_n is the maximal one.

Definition 35 Let $\leq_B$ be the relation on $\mathcal{T}_n$ generated transitively by the following relations:

1. If $t^{i_0} <_B w^{i_0} \in \mathcal{T}_{n_{i_0}}$, then $\bigvee(t^0, \dots, t^{i_0}, \dots, t^k) <_B \bigvee(t^0, \dots, w^{i_0}, \dots, t^k)$ in $\mathcal{T}_{n_0 + \dots + n_k + k - 1}$,

2. If $t = \bigvee(t^0, \dots, t^k) \in \mathcal{T}_n$ and $w = \bigvee(w^0, \dots, w^h) \in \mathcal{T}_m$, then


$$\bigvee(t, w^0, \dots, w^h) <_B \bigvee(t^0, \dots, t^k, w^0, \dots, w^h)$$

3. If $t = \bigvee(t^0, \dots, t^k) \in \mathcal{T}_n$, then

$$t <_B \bigvee(t^0, \dots, t^j, \bigvee(t^{j+1}, \dots, t^k)),$$

for $1 \leq j < k$.

Example 36 On $\mathcal{T}_2$, we have



Proposition 37 The pair $(\mathcal{T}_n, \leq_B)$ is a partially ordered set, for all $n \geq 1$.

Proof. We need to check that, given a tree t in $\mathcal{T}_n$, it does not exist a sequence of trees $t = t_0, t_1, \dots, t_m = t$ such that t_{i+1} is obtained from t_i applying one of the relations 1, 2 or 3 of Definition 35.

We proceed by induction on n . For $n = 1$ or 2 the result is immediate. Suppose now that $|t| > 2$, there exist trees $t^0, \dots, t^k$ such that $t = \bigvee(t^0, \dots, t^k)$. Note that, if t_i is obtained from t_{i-1} by an operation of type 1 or 3, then $|t_i^1| = |t_{i-1}^1|$, while applying operation of type 2, we get $|t_i^1| < |t_{i-1}^1|$. So, it there exists a sequence $t = t_0, t_1, \dots, t_m = t$ such that t_{i+1} it must be obtained applying only operations of types 1 or 3.

Now, suppose $t_j = \bigvee(t_j^0, \dots, t_j^{k_j})$, for $0 \leq j \leq m$. If t_j is obtained from t_{j-1} by an operation of type 1, then $|t_j^{r_j}| = |t_{j-1}^{r_{j-1}}|$, and if it is obtained from t_{j-1} by an operation of type 2, then $|t_j^{r_j}| > |t_{j-1}^{r_{j-1}}|$. We may conclude then that the sequence $t = t_0, t_1, \dots, t_m = t$ must be obtained applying only operations of type 1.

In this case, it must exist at least one $1 \leq k \leq m$ and a sequence $t^k = t_0^k <_B \dots <_B t_m^k = t^k$. But, by induction hypothesis, we know that it is false. So, the relation $\leq_B$ is a well defined order on $\mathcal{T}_n$. $\diamond$

It is immediate to check that $\leq_B$ induces an order on the subset $\mathcal{Y}_n$ of $\mathcal{T}_n$. This order is the order defined in [LRo].

Remark 38 Let $t = \bigvee(t^0, \dots, t^k)$ and $z = \bigvee(z^0, \dots, z^k)$ in $\mathcal{T}_n$, such that $|t^i| = |z^i|$, for $0 \leq i \leq k$.

1. If $t \subseteq z$ (respectively, $t \leq_B z$), then $t^i \subseteq z^i$ (respectively $t^i \leq_B z^i$), for all $0 \leq i \leq k$.

2. If $w \in \mathcal{T}_n$ is such that $t \subseteq w \subseteq z$ (respectively $t \leq_B w \leq_B z$), then
 $w = \bigvee(w^0, \dots, w^k)$ with $t^i \subseteq w^i \subseteq z^i$ (respectively $t^i \leq_B w^i \leq_B z^i$), for
 $0 \leq i \leq k$.

3.2 Poset morphism from $\mathcal{P}_\infty$ to $\mathcal{T}_\infty$

Definition 39 Let us define a map $\Gamma : \mathcal{P}_\infty \longrightarrow \mathcal{T}_\infty$ as follows:

1. $\Gamma((0)) = \downarrow$, where $\downarrow$ is the tree with one leaf and without vertex.
2. Let $\gamma = \bigvee_\omega(\gamma_0, \dots, \gamma_k)$ be an element of $\mathcal{P}_n$. The element $\Gamma(\gamma)$ is the tree:

$$\Gamma(\gamma) := \bigvee(\Gamma(\gamma_0), \dots, \Gamma(\gamma_k)).$$

Remark 40 1. It is immediate to see that $\Gamma((1, \dots, 1)) = \varsigma_n$, for $n \geq 1$.

2. An easy recursive argument shows that Γ is surjective.
3. Let $\gamma_0, \dots, \gamma_k$ be a family of elements in $\mathcal{P}_\infty$, with $\gamma_i \in \mathcal{P}_{n_i, r_i}$ and $r = \sum_{i=0}^k r_i$, and let ω be an element in $SH(r_0, \dots, r_k)$. The images of the elements $\bigvee_\omega(\gamma_0, \dots, \gamma_k)$ and $\bigvee_{1_r}(\gamma_0, \dots, \gamma_k)$ under Γ are equal.
4. The image of the subset $\mathcal{P}_{n,r}$ of $\mathcal{P}_n$ under Γ is the subset $\mathcal{T}_{n,r} := \{t \in \mathcal{T}_n : t \text{ has } n - r \text{ vertices}\}$.

Theorem 41 The map $\Gamma : \mathcal{P}_\infty \rightarrow \mathcal{T}_\infty$ verifies that:

1. If $\gamma \subseteq \delta$ in $\mathcal{P}_n$, then $\Gamma(\gamma) \subseteq \Gamma(\delta)$ in $\mathcal{T}_n$.
2. If $\gamma \leq_B \delta$ in $\mathcal{P}_n$, then $\Gamma(\gamma) \leq_B \Gamma(\delta)$ in $\mathcal{T}_n$.

Proof. In this proof, given a sequence of integers $r_0, \dots, r_k$, we shall denote by R_l the sum $R_l := r_0 + \dots + r_l$, for $0 \leq l \leq k$.

1. We know that $\gamma \subseteq \delta$ if, and only if, there exists a non-decreasing map ρ such that $\delta = \rho \circ \gamma$. Since ρ is a product of t_i 's, it suffices to prove the result for $\rho = t_j$.

The result is immediate for $n = 0, 1, 2$. For $n \geq 3$, suppose $\gamma = \bigvee_\omega(\gamma_0, \dots, \gamma_k)$, with $\gamma_i \in \mathcal{P}_{n_i, r_i}$. For $j < \sum_{i=0}^k r_i$, we have that:

- (a) If $t_j \circ \omega \in SH(r_0, \dots, r_k)$, then $\delta = \bigvee_{t_j \circ \omega}(\gamma_0, \dots, \gamma_k)$ and $\Gamma(\delta) = \Gamma(\gamma)$.
- (b) If $t_j \circ \omega \notin SH(r_0, \dots, r_k)$, then Lemma 21 implies that $\delta = \bigvee_{\omega_j}(\gamma_0^j, \dots, \gamma_k^j)$, where $\gamma_i^j = \gamma_i$ or $\gamma_i^j = t_{j_i} \circ \gamma_i$. So, $\gamma_i \subseteq \gamma_i^j$, for $0 \leq i \leq k$.
By recursive hypothesis $\Gamma(\gamma_i) \subseteq \Gamma(\gamma_i^j)$, and $\Gamma(\delta) = \bigvee(\Gamma(\gamma_0^j), \dots, \Gamma(\gamma_k^j))$.
So, we get that $\Gamma(\gamma) \subseteq \Gamma(\delta)$.

For $j = \sum_{i=0}^k r_i = r - 1$, the element δ is obtained as a wedge of two types of elements:

$$\begin{cases} \gamma_i & \text{for } \omega(R_i) < r - 1, \\ \gamma_{l,i} & \text{for } \omega(R_i) = r - 1 \text{ and } \gamma_i = \bigvee_{\omega^i}(\gamma_{0,i}, \dots, \gamma_{k_i,i}). \end{cases}$$

So, $\Gamma(\delta)$ is the wedge of the elements:

$$\begin{cases} \Gamma(\gamma_i) & \text{for } \omega(R_i) < r - 1, \\ \Gamma(\gamma_{l,i}) & \text{for } \omega(R_i) = r - 1 \text{ and } \gamma_i = \bigvee_{\omega^i}(\gamma_{0,i}, \dots, \gamma_{k_i,i}). \end{cases}$$

The argument above implies that $\Gamma(\delta)$ is obtained from $\Gamma(\gamma)$ by contracting all the edges which joint the root of $\Gamma(\gamma_i)$ to the root of $\Gamma(\gamma)$, for all $0 \leq i \leq k$ such that $\omega(R_i) = r - 1$. It implies that $\Gamma(\gamma) \subseteq \Gamma(\delta)$.

2. For the weak Bruhat order the result is immediate for $n \leq 2$. For $n \geq 3$, suppose $\gamma = \bigvee_{\omega}(\gamma_0, \dots, \gamma_k)$, and $\delta = \bigvee_v(\delta_0, \dots, \delta_h)$, with $\gamma_i \in \mathcal{P}_{n_i, r_i}$ and $\delta_i \in \mathcal{P}_{m_i, s_i}$.

a) If $\delta = t_j \circ \gamma$, with $\gamma^{-1}(j) < \gamma^{-1}(j + 1)$, then we have that:

- (i) For $j < r - 1$ and $t_j \circ \omega \in SH(r_0, \dots, r_k)$, it holds that $\Gamma(\delta) = \Gamma(\gamma)$.
- (ii) For $j < r - 1$ and $t_j \circ \omega \notin SH(r_0, \dots, r_k)$, Lemma 21 states that $\delta = \bigvee_{\omega_j}(\gamma_0^j, \dots, \gamma_k^j)$, with

$$\begin{cases} \gamma_i^j = \gamma_i & \text{when } \{j, j + 1\} \not\subseteq \omega(\{R_{i-1} + 1, \dots, R_i\}), \\ \gamma_i^j = t_{j_i} \circ \gamma_i & \text{when } \{j, j + 1\} \subseteq \omega(\{R_{i-1} + 1, \dots, R_i\}). \end{cases}$$

The fact that $\gamma^{-1}(j) < \gamma^{-1}(j + 1)$ implies that there exists at most one i_0 such that $\gamma_{i_0}^j \neq \gamma_{i_0}$ and $\gamma_{i_0}^{-1}(j_{i_0}) < \gamma_{i_0}^{-1}(j_{i_0} + 1)$. So,

$$\Gamma(\delta) = \bigvee(\Gamma(\gamma_0), \dots, \Gamma(t_{j_{i_0}} \circ \gamma_{i_0}), \dots, \Gamma(\gamma_k)),$$

and, by recursive hypothesis, $\Gamma(\gamma_{i_0}) \leq_B \Gamma(t_{j_{i_0}} \circ \gamma_{i_0})$, which implies that

$$\Gamma(\gamma) \leq_B \Gamma(\delta).$$

(iii) For $j = r - 1$, we get that $\gamma^{-1}(r - 1) \subseteq \{1, \dots, r_0\}$. It holds that

$$\Gamma(\gamma) = \bigvee(\Gamma(\gamma_0), \dots, \Gamma(\gamma_k))$$

and that $\Gamma(\delta)$ is the wedge of $\Gamma(\gamma_{0,0}), \dots, \Gamma(\gamma_{k_0,0}), \Gamma(\gamma_1), \dots, \Gamma(\gamma_k)$, with

$$\Gamma(\gamma_0) = \bigvee(\Gamma(\gamma_{0,0}), \dots, \Gamma(\gamma_{k_0,0})).$$

So, we get that $\Gamma(\gamma) \leq_B \Gamma(\delta)$.

b) The proof of the case $t_j \circ \delta = \gamma$, with $\delta^{-1}(j) > \delta^{-1}(j + 1)$, can be obtained following the same arguments that we use to prove a). $\diamond$

In order to see that the orders $\subseteq$ and $\leq_B$ of $\mathcal{P}_\infty$ induce the orders $\subseteq$ and $\leq_B$ on $\mathcal{T}_\infty$, we need to prove some additional results.

Remark 42 *Let γ, ω and δ be elements of $\mathcal{P}_m$ such that $\Gamma(\gamma) = \Gamma(\delta)$ and $\gamma \subseteq \omega \subseteq \delta$. The theorem above implies that $\Gamma(\omega) = \Gamma(\gamma) = \Gamma(\delta)$.*

Let $\leq$ be the relation on $\mathcal{T}_\infty$ given by $t \leq z$ if and only if there exist $\gamma \in \Gamma^{-1}(t)$ and $\delta \in \Gamma^{-1}(z)$ such that $\gamma \subseteq \delta$.

Proposition 43 *The relation $\leq$ defines a partial order on $\mathcal{T}_\infty$ which coincides with $\subseteq$.*

Proof. It results easy to check that $\leq$ is reflexive and transitive. Lemma 21 states that if $\delta = \rho \circ \gamma$, with ρ a non decreasing element in $\mathcal{P}_r$, then either $\Gamma(\delta) = \Gamma(\gamma)$ or $\Gamma(\delta)$ is obtained from $\Gamma(\gamma)$ by contracting some internal edges. This fact implies the antisymmetry of $\leq$.

If $t \leq z$, then there exist $\gamma \in \Gamma^{-1}(t)$ and $\delta \in \Gamma^{-1}(z)$ with $\gamma \subseteq \delta$. By Theorem 41, it holds that $t = \Gamma(\gamma) \subseteq \Gamma(\delta) = z$.

If $t \subseteq z$, then it becomes clear that $t \leq z$ for $|t| = |z| \leq 2$. For $|t| = |z| > 2$, suppose that z is obtained from $t = \bigvee(t^0, \dots, t^k)$ by contracting an internal edge.

1. If $z = \bigvee(t^0, \dots, z^j, \dots, t^k)$, with $t^j \subseteq z^j$, then by recursive hypothesis there exist $\gamma_j \in \Gamma^{-1}(t^j)$ and $\delta_j \in \Gamma^{-1}(z^j)$ such that $\delta_j = \rho_j \circ \gamma_j$, with ρ_j a non-decreasing map in $\mathcal{P}_\infty$.

Choose an element $\gamma_i \in \Gamma^{-1}(t^i)$, for $i \neq j$. The element $\gamma := \bigvee_{1_{r-1}}(\gamma_0, \dots, \gamma_k)$ belongs to $\Gamma^{-1}(t)$. Consider $\delta := \bigvee_{1_{r-1-r_j+s_j}}(\gamma_0, \dots, \delta_j, \dots, \gamma_k)$, we get that $\delta \in \Gamma^{-1}(z)$ and $\gamma \subseteq \delta$, which implies $t \leq z$.

2. For $z = \bigvee(t^0, \dots, t^{0,j}, \dots, t^{n_j,j}, \dots, t^k)$, with $t^j = \bigvee(t^{0,j}, \dots, t^{n_j,j})$, choose elements γ_i be in $\Gamma^{-1}(t^i)$, for $i \neq j$, and $\gamma_{l,j} \in \Gamma^{-1}(t^{l,j})$, for $0 \leq l \leq n_j$.

The element $\gamma_j := \bigvee_{1_{r_j-1}}(\gamma_{0,j}, \dots, \gamma_{n_j,j})$ belongs to $\Gamma^{-1}(\gamma_j)$, $\gamma := \bigvee_{1_{r-1}}(\gamma_0, \dots, \gamma_k)$ is in $\Gamma^{-1}(t)$, and $\delta := \bigvee_{1_{r-1-r_j+\sum r_{l,j}}}(\gamma_0, \dots, \gamma_{0,j}, \dots, \gamma_{n_j,j}, \dots, \gamma_k)$ belongs to $\Gamma^{-1}(z)$. Since $\gamma \subseteq \delta$ we may assert that $t \leq z$. $\diamond$

Definition 44 Given a tree $t \in \mathcal{T}_m$, we define the elements $Min(t)$ and $Max(t)$ in $\mathcal{P}_m$ recursively, as follows:

1. $Min(\downarrow) := (0) =: Max(\downarrow)$.
2. For $t = \bigvee(t^0, \dots, t^k)$, with $t^i \in \mathcal{T}_{n_i}$ and $n = \sum_{i=0}^k n_i$, the degree of t is $n + k$. Define

$$Min(t) := \bigvee_{1_{r-1}}(Min(t^0), \dots, Min(t^k)),$$

$$Max(t) := \bigvee_{\xi_{s_0, \dots, s_k}}(Max(t^0), \dots, Max(t^k)),$$

where $\xi_{s_0, \dots, s_k}$ is the biggest element of $SH(s_0, \dots, s_k)$ for the weak Bruhat order, $Min(t^i) \in \mathcal{P}_{n_i, r_i}$, $Max(t^i) \in \mathcal{P}_{n_i, s_i}$ and $\sum_{i=0}^k r_i = r - 1$.

Proposition 45 Let $t \in \mathcal{T}_m$ be a tree, the inverse image under Γ of t is an interval for the weak Bruhat order. Moreover, it verifies

$$\Gamma^{-1}(t) = \{\delta \in \mathcal{P}_m \text{ such that } Min(t) \leq_B \delta \leq_B Max(t)\}.$$

Proof. The assertion is obviously true for $m = 0, 1, 2$. For $m \geq 3$, let $t \in \mathcal{T}_m$ with $t = \bigvee(t^0, \dots, t^k)$. By recursive hypothesis, we know that

$$\Gamma^{-1}(t^i) = \{\delta_i \in \mathcal{P}_{n_i} \text{ such that } Min(t^i) \leq_B \delta_i \leq_B Max(t^i)\}.$$

Suppose $\gamma = \bigvee_{\omega}(\gamma_0, \dots, \gamma_k)$ is such that $\Gamma(\gamma) = t$. We have that $\Gamma(\gamma_i) = t^i$, for $0 \leq i \leq k$. By Corollary 27 and Lemma 28 we obtain that:

$$Min(t) = \bigvee_{1_{r-1}}(Min(t^0), \dots, Min(t^k)) \leq_B \bigvee_{1_{r'-1}}(\gamma_0, \dots, \gamma_k) \leq_B \gamma,$$

and, there exists $\omega' \leq_B \xi_{s_0, \dots, s_k}$ such that:

$$\gamma \leq_B \bigvee_{\omega'} (Max(t^0), \dots, Max(t^k)) \leq_B \bigvee_{\xi_{s_0, \dots, s_k}} (Max(t^0), \dots, Max(t^k)) = Max(t).$$

Conversely, if $Min(t) \leq_B \gamma \leq_B Max(t)$, then Theorem 41 states that

$$t = \Gamma(Min(t)) \leq_B \Gamma(\gamma) \leq_B \Gamma(Max(t)) = t. \diamond$$

So, the weak Bruhat order of $\mathcal{P}_\infty$ induces a partial order on $\mathcal{T}_\infty$.

Proposition 46 *The weak Bruhat order $\leq_B$ of $\mathcal{P}_\infty$ induces the weak Bruhat order on $\mathcal{T}_\infty$.*

Proof. Using that Γ preserves the weak Bruhat order, it is easily seen that $t \leq z$ for the order induced by the weak Bruhat order of $\mathcal{P}_\infty$ implies that $t \leq_B z$.

It is immediate to check that $t \leq_B z$ implies $t \leq z$ for the induced order, when $|t| = |z| \leq 2$. For $|t| \geq 3$, we have to consider the following three cases:

1. If $t = \bigvee(t^0, \dots, t^k)$ and $z = \bigvee(z^0, \dots, z^k)$, with $t^i \leq_B z^i$ for all i , then by recursive hypothesis we get that $t^i \leq z^i$ for the induced order.

So, there exist two families $\gamma_0, \dots, \gamma_k$ and $\delta_0, \dots, \delta_k$ of element in $\mathcal{P}_\infty$ such that $\gamma_i \leq_B \delta_i$, $\Gamma(\gamma_i) = t^i$ and $\Gamma(\delta_i) = z^i$. Since $t = \Gamma(\bigvee_{1_{r-1}}(\gamma_0, \dots, \gamma_k))$, $z = \Gamma(\bigvee_{1_{s-1}}(\delta_0, \dots, \delta_k))$ and $\bigvee_{1_{r-1}}(\gamma_0, \dots, \gamma_k) \leq_B \bigvee_{1_{s-1}}(\delta_0, \dots, \delta_k)$, we get that $t \leq z$ for the induced order.

2. If $t = \bigvee(t^0, \dots, t^k)$ and $z = \bigvee(t^{0,0}, \dots, t^{k_0,0}, t^1, \dots, t^k)$, for $t^0 = \bigvee(t^{0,0}, \dots, t^{k_0,0})$, then there exist $\gamma_{0,0}, \dots, \gamma_{k_0,0}, \gamma_1, \dots, \gamma_k$ in $\mathcal{P}_\infty$ such that $\Gamma(\gamma_{l,0}) = t^{l,0}$, for $0 \leq l \leq k_0$, and $\Gamma(\gamma_i) = t^i$, for $1 \leq i \leq k$.

Consider the elements $\gamma_0 := \bigvee_{1_{n_0}}(\gamma_{0,0}, \dots, \gamma_{k_0,0})$, $\gamma := \bigvee_{\xi_{r_0, r-1-r_0}}(\gamma_0, \dots, \gamma_k)$ and $\delta := \bigvee_{1_{s-1}}(\gamma_{0,0}, \dots, \gamma_{k_0,0}, \gamma_1, \dots, \gamma_k)$, where $\gamma_i \in \mathcal{P}_{n_i, r_i}$, $\gamma_{l,0} \in \mathcal{P}_{n_{l,0}, r_{l,0}}$, $\sum_{i=0}^k r_i = r-1$ and $\sum_{l=0}^{k_0} r_{l,0} + \sum_{i=1}^k r_i = s-1$. It holds that $\Gamma(\gamma) = t$, $\Gamma(\delta) = z$, $\gamma^{-1}(r-1) < \gamma^{-1}(r)$ and $\delta = t_{r-1} \circ \gamma$, which implies $t \leq z$ for the induced order.

3. The proof for the case $t = \bigvee(z^0, \dots, z^k, z^{0,k}, \dots, z^{0,h_k})$ and $z = \bigvee(z^0, \dots, z^k)$ is obtained in the same way that the one of case 2. $\diamond$

3.3 Free dendriform trialgebra structure on $k[\mathcal{T}_\infty]$

Note that the operations $\succ$, $\cdot$ and $\prec$ defined in Section 2 induce operations on the k -vector space $k[\mathcal{T}_\infty]$, spanned by the set of planar trees. The operations are defined as follows:

1. $t \succ z = \sum w$, where the sum is taken over the set $\{w \in \mathcal{T}_\infty \text{ such that there exist } \gamma \in \Gamma^{-1}(t), \delta \in \Gamma^{-1}(z) \text{ and } \sigma \in SH^\succ(r, s) \text{ such that } \sigma \circ (\gamma \times \delta) \in \Gamma^{-1}(w)\}$.
2. $t \cdot z = \sum w$, where the sum is taken over the set $\{w \in \mathcal{T}_\infty \text{ such that there exist } \gamma \in \Gamma^{-1}(t), \delta \in \Gamma^{-1}(z) \text{ and } \sigma \in SH^\bullet(r, s) \text{ such that } \sigma \circ (\gamma \times \delta) \in \Gamma^{-1}(w)\}$.
3. $t \prec z = \sum w$, where the sum is taken over the set $\{w \in \mathcal{T}_\infty \text{ such that there exist } \gamma \in \Gamma^{-1}(t), \delta \in \Gamma^{-1}(z) \text{ and } \sigma \in SH^\prec(r, s) \text{ such that } \sigma \circ (\gamma \times \delta) \in \Gamma^{-1}(w)\}$.

Remark 47 Given a partition $n_0, \dots, n_k$ of n and $1 \leq j \leq n$, we denote by $SH(n_0, \dots, n_k)_r$ the set of elements $\omega \in SH(n_0, \dots, n_k) \cap \mathcal{P}_{n,r}$. An easy calculation shows that the following equalities hold:

$$SH(r, s_0, \dots, s_h) = \bigcup_j SH(r, j) \circ (1_r \times SH(s_0, \dots, s_h)) = \bigcup_j SH(j, s_1, \dots, s_h) \circ (SH(r, s_0) \times 1_{s-s_0-1}), \quad (1)$$

$$\bigcup_{i,j} SH(i, j) \circ (SH(r_0, \dots, r_k)_i \times SH(s_0, \dots, s_h)_j) = \bigcup_i SH(r_0, \dots, r_{k-1}, i, s_1, \dots, s_h) \circ (1_{r-r_k-1} \times SH(r_k, s_0) \times 1_{s-s_0-1}), \quad (2)$$

$$SH(r_0, \dots, r_k, s) = \bigcup_j SH(j, s) \circ (SH(r_0, \dots, r_k) \times 1_s) = \bigcup_j SH(r_0, \dots, r_{k-1}, j) \circ (1_{r-r_k-1} \times SH(r_k, s)), \quad (3)$$

where $r - 1 = \sum_{i=0}^k r_i$ and $s - 1 = \sum_{i=0}^h s_i$.

Theorem 48 The operations $\succ$, $\cdot$ and $\prec$ induced on $k[\mathcal{T}_\infty]$ by the dendriform trialgebra structure of $k[\mathcal{P}_\infty]$ verify the following equalities:

1. $t * \downarrow = t = \downarrow * t$, for all $t \in \mathcal{T}_\infty$,
2. $t \succ z = \bigvee (t * z^0, z^1, \dots, z^h)$,

$$3. t \cdot z = \bigvee(t^0, \dots, t^{k-1}, t^k * z^0, z^1, \dots, z^h),$$

$$4. t \prec z = \bigvee(t^0, \dots, t^{k-1}, t^k * z),$$

for $t = \bigvee(t^0, \dots, t^k)$ and $z = \bigvee(z^0, \dots, z^h)$.

Proof. 1 is obvious.

The proofs of 2, 3 and 4 use similar arguments, applying the Remark above. We give the detailed proof of 3 (which seems to be just a bit more difficult), the other ones may be obtained following analogous steps.

3. We have that $SH^\bullet(j, l) = (SH(j-1, l-1) \times 1_1) \circ z(j-1, l-1)$. A tree w appears with coefficient 1 in $t \cdot z$ if there exist $\{\gamma_i \in \mathcal{P}_{n_i, r_i}\}_{0 \leq i \leq k}$, $\omega \in SH(r_0, \dots, r_k)_{j-1}$, $\{\delta_i \in \mathcal{P}_{m_i, s_i}\}_{0 \leq i \leq h}$, $v \in SH(s_0, \dots, s_h)_{l-1}$ and $\sigma \in SH(j-1, l-1)$ such that the element $\sigma \circ (\gamma \times \delta)$ belongs to $\Gamma^{-1}(w)$, with $\sigma = (\sigma_1 \times 1_1)$, $\gamma = \bigvee_\omega(\gamma_0 \times \dots \times \gamma_k)$ and $\delta = \bigvee_v(\delta_0 \times \dots \times \delta_h)$. Using Remark 47 (2), we get that

$$\begin{aligned} \sigma \circ (\gamma \times \delta) &= (\sigma_1 \times 1_1) \circ z(j-1, l-1) \circ \\ &\quad ((\omega \circ (\gamma_0 \times \dots \times \gamma_k)) \times 1_1 \times (v \circ (\delta_0 \times \dots \times \delta_h)) \times 1_1) \circ \\ &\quad (z(n_0, \dots, n_k) \times z(m_0, \dots, m_h)) = \\ &= (\sigma_1 \times 1_1) \circ ((\omega \circ (\gamma_0 \times \dots \times \gamma_k)) \times v \circ (\delta_0 \times \dots \times \delta_h) \times 1_1) \\ &\quad \circ z(n_0, \dots, n_k, m_0, \dots, m_h) = \\ &= \tau \circ (1_{r-r_0-1} \times \zeta \times 1_{s-s_0-1}) \circ \\ &\quad (\gamma_0 \times \dots \times \gamma_k \times \delta_0 \times \dots \times \delta_h \times 1_1) \circ z(n_0, \dots, n_k, m_0, \dots, m_h) = \\ &\quad \bigvee_\tau (\gamma_0, \dots, \gamma_{k-1}, \zeta \circ (\gamma_k \times \delta_0), \delta_1, \dots, \delta_h), \end{aligned}$$

for unique elements $\zeta \in SH(r_k, s_0)_i$ and $\tau \in SH(r_0, \dots, r_{k-1}, i, s_1, \dots, s_h)$. So, w appears in $\bigvee(t^0, \dots, t^{k-1}, t^k * z^0, z^1, \dots, z^h)$.

Looking at the same equalities from the lower line to the upper one, we show that any tree w which appears in $\bigvee(t^0, \dots, t^{k-1}, t^k * z^0, z^1, \dots, z^h)$ with coefficient 1 appears with the same coefficient in $t \cdot z$, too. $\diamond$

The following result is a consequence of the Theorem above and the description of the free dendriform trialgebra given in [LR4].

Corollary 49 *The dendriform trialgebra $(k[\mathcal{T}_\infty], \succ, \cdot, \prec)$ is the free dendriform trialgebra spanned by one element.*

Let us describe now the operations of the free dendriform algebra $k[\mathcal{T}_\infty]$ in terms of the weak Bruhat order of $\mathcal{T}_\infty$.

Given a tree $t \in \mathcal{T}_n$, we numerate its leaves from left to right by $0, 1, \dots, n$.

Notation 50 Let $t \in \mathcal{T}_n$ and $z \in \mathcal{T}_m$ be two planar trees.

1. For any $0 \leq j \leq m$, the grafting $t \circ_j z$ of t on the j -th leaf of z is the tree obtained by joining the root of t to the leaf j of z .
2. We denote by t/z the tree $t \circ_0 z$, and by $t \setminus z$ the tree $z \circ_n t$.

Proposition 51 Let $t \in \mathcal{T}_n$ and $z \in \mathcal{T}_m$. Given elements $\gamma \in \Gamma^{-1}(t) \cap \mathcal{P}_{n,r}$ and $\delta \in \Gamma^{-1}(z) \cap \mathcal{P}_{m,s}$, it holds that:

1. $\Gamma(\gamma \times \delta) = t/z$,
2. $\Gamma(\xi_{r,s} \circ (\gamma \times \delta)) = t \setminus z$.

Proof.

1. If $z = \downarrow$, then $\gamma \times (0) = \gamma$ and $\Gamma(\gamma) = t = t/\downarrow$.

If $m \geq 1$, then $\delta = \bigvee_v (\delta_0, \dots, \delta_h)$, for a family $\delta_i \in \Gamma^{-1}(z^i)$, with $0 \leq i \leq h$, and $v \in SH(s_0, \dots, s_h)$. We have that:

$$\begin{aligned} \gamma \times \delta &= (1_r \times v) \circ (\gamma \times \delta_0 \times \dots \times \delta_h \times 1_1) \circ (1_n \times z(m_0, \dots, m_h)) = \\ &= (1_r \times v) \circ ((\gamma \times \delta_0) \times \delta_1 \times \dots \times \delta_h \times 1_1) \circ z(n + m_0, m_1, \dots, m_h) = \\ &= \bigvee_{1_r \times v} (\gamma \times \delta_0, \delta_1, \dots, \delta_h). \end{aligned}$$

The recursive hypothesis states that $\Gamma(\gamma \times \delta_0) = t/z^0$, which implies that $\Gamma(\gamma \times \delta) = \bigvee (t/z^0, z^1, \dots, z^h) = t/z$.

2. If $t = \downarrow$, then $\xi_{0,r} \circ ((0) \times \delta) = \delta$ and $\Gamma(\delta) = z = \downarrow \setminus z$.

If $n \geq 1$, then $\gamma = \bigvee_\omega (\gamma_0, \dots, \gamma_k)$, with $\gamma_i \in \Gamma^{-1}(t^i)$ and $\omega \in SH(r_0, \dots, r_k)_r$, with $\sum_{i=0}^k r_i = r' - 1$. It holds that

$$\begin{aligned} \xi_{r,s} \circ (\gamma \times \delta) &= \xi_{r,s} \circ (\omega \times 1_{s+1}) \circ (\gamma_0 \times \dots \times \gamma_k \times 1_1 \times \delta) \circ (z(r_0, \dots, r_k) \times 1_m) = \\ &= (\omega(1)+s, \dots, \omega(r'-1)+s, r+s, 1, \dots, s) \circ (\gamma_0 \times \dots \times \gamma_k \times 1_1 \times \delta) \circ (z(r_0, \dots, r_k) \times 1_m) = \\ &= ((\xi_{r-1,s} \circ (\omega \times 1_s) \circ (\gamma_0 \times \dots \times \gamma_{k-1} \times (\gamma_k \times \delta))) \times 1_1) \circ z(r_0, \dots, r_{k-1}, r_k + s) = \\ &= (v \times 1_1) \circ (\gamma_0 \times \dots \times \gamma_{k-1} \times (\xi_{r_k,s} \circ (\gamma_k \times \delta))) \times 1_1 \circ z(r_0, \dots, r_{k-1}, r_k + s) = \\ &= \bigvee_v (\gamma_0, \dots, \gamma_{k-1}, (\xi_{r_k,s} \circ (\gamma_k \times \delta))), \end{aligned}$$

where $v :=$

$(\omega(1)+s, \dots, \omega(r_0+\dots+r_{k-1})+s, 1, \dots, s, \omega(r_0+\dots+r_{k-1}+1)+s, \dots, \omega(r'-1)+s),$
 belongs to $SH(r_0, \dots, r_{k-1}, r_k + s).$

By recursive hypothesis, $\Gamma(\xi_{r_k, s} \circ (\gamma_k \times \delta)) = t^k \setminus z.$

So, we get that $\Gamma(\xi_{r, s} \circ (\gamma \times \delta)) = \bigvee(t^0, \dots, t^{k-1}, t^k \setminus z) = t \setminus z. \quad \diamond$

Theorem 48 and Proposition 51 imply the following result:

Corollary 52 *The operations $\succ, \cdot, \prec$ and $*$ of the free dendriform trialgebra $k[\mathcal{T}_\infty]$ are defined in terms of the weak Bruhat order and the products $/$ and $\setminus$ by the following conditions:*

1. $t \succ z = \sum_{t/z \leq_B w \leq_B \bigvee(t \setminus z^0, z^1, \dots, z^k)} w,$
2. $t \cdot z = \sum_{\bigvee(t^0, \dots, t^{k-1}, t^k / z^0, z^1, \dots, z^h) \leq_B w \leq_B \bigvee(t^0, \dots, t^{k-1}, t^k \setminus z^0, z^1, \dots, z^h)} w,$
3. $t \prec w = \sum_{\bigvee(t^0, \dots, t^{k-1}, t^k / z) \leq_B w \leq_B t \setminus z} w,$
4. $t * z = \sum_{t/z \leq_B w \leq_B t \setminus z} w.$

Final comment Given a Coxeter system (W, S) there exists another order on the set $\mathcal{P}^{(W, S)}$ which extends the weak Bruhat order of W . Consider the transitive relation $\leq_C$ spanned by:

$$W_J \circ w \leq_C W_J \circ (\alpha_{J, s}^{-1} \circ w), \text{ for } J \subseteq S, s \in S \setminus J \text{ and } w \in X_{J \cup \{s\}}^{-1},$$

where W_J , X_K and $\alpha_{J, s}$ are the elements defined in Section 1.

Observe that if $x \leq_C y$, then $x \leq_B y$. So, the weak Bruhat order is weaker than $\leq_C$.

This new order preserves the graduation of $\mathcal{P}^{(W, S)}$, that is two elements x and y of $\mathcal{P}^{(W, S)}$ are comparable for $\leq_C$ only if $x, y \in \mathcal{P}_r^{(W, S)}$, for some $0 \leq r \leq |S|$.

Mimicking the results of Section 3 it is possible to see that the order $\leq_C$ of $\mathcal{P}_\infty$ induces, via the map Γ , a partial order on $\mathcal{T}_\infty$.

The order $\leq_C$ on $\mathcal{P}_n$ is the transitive relation generated by:

$$\omega \leq_C s_i \circ \omega, \quad \text{for } \omega^{-1}(i) \leq \omega^{-1}(i+1),$$

where $\omega^{-1}(j)$ denotes the set of elements l of $\{1, \dots, n\}$ such that $\omega(l) = j$.

On $\mathcal{T}_n$, the order $\leq_C$ is transitively spanned by the relations:

1. If $t^{i_0} \leq_C w^{i_0} \in \mathcal{T}_{n_{i_0}}$, then $\bigvee(t^0, \dots, t^{i_0}, \dots, t^k) \leq_C \bigvee(t^0, \dots, w^{i_0}, \dots, t^k)$ in $\mathcal{T}_{n_0 + \dots + n_k + k - 1}$,
2. If $t = \bigvee(t^0, \dots, t^k) \in \mathcal{T}_n$ and $w = \bigvee(w^0, \dots, w^h) \in \mathcal{T}_m$, then

$$\bigvee(t, w^0, \dots, w^h) \leq_C \bigvee(t^0, \dots, t^k, w).$$

In this case, the order $\leq_C$ also permits to construct associative algebra structures on $k[\mathcal{P}_\infty]$ and $k[\mathcal{T}_\infty]$. These structures coincide with the associative algebra structures defined on $k[\mathcal{P}_\infty]$ and $k[\mathcal{T}_\infty]$ by F. Chapoton in [Ch].

References

- [BBHT] F. Bergeron, N. Bergeron, R. Howlett and R. Taylor. *A decomposition of the descent algebra of a finite Coxeter group*, J. Algebraic Combin. 1 (1992), no. 1, 23–44.
- [Bo] N. Bourbaki, *Groupes et algèbres de Lie*, Ch. 4-6, Herman- Paris (1968); Masson-Paris (1981).
- [Ch] F. Chapoton, *Algèbres de Hopf des permutoèdres, associaèdres et hypercubes*, Advances in Math. 150 (2000), 264-275.
- [DHT] G. Duchamp, F. Hivert and J.-Y. Thibon, *Noncommutative symmetric functions VI: free quasi-symmetric functions and related algebras*, Intern. J. of Algebra and Computation 2 (2002) 671-717.
- [GK] V. Ginzburg y M. Kapranov. *Koszul duality for operads*, Duke Math. J., 76:203-272, 1994.
- [GKLLR] I.M. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V. Retakh, J.-Y. Thibon, *Noncommutative symmetric functions*, Adv. Math. 112 (1995), no. 2, 218-348.
- [HNT] F. Hivert, J.-C. Novelli et J.-Y. Thibon, *Un analogue du monoïde plaxique pour les arbres binaires de recherche*, C. R. Acad. Sci. Paris, 335 (2002) 577-580.
- [Hu] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge studies in advanced mathematics 29, Cambridge Univ. Press, Reprint edition (1993).

- [L1] J.-L. Loday. *Arithmetree*, J. of Algebra 258 (2002) 275-309.
- [L2] J.-L. Loday. *Dialgebras* , in Dialgebras and related operads, Springer Lecture Notes in Math. 1763 (2002), 7-66.
- [LR1] J.-L. Loday, M. Ronco, *The Hopf algebra of planar binary trees* , Advances in Maths. 139, 293-309 (1998).
- [LR2] J.-L. Loday, M. Ronco, *Order structure and the algebra of permutations and of planar binary trees* , J. of Algebraic Combinatorics 15 (2002), 253-270.
- [LR3] J.-L. Loday, M. Ronco, *Une dualité entre simplexes standards et polytopes de Stasheff* , C. R. Acad. Sci. Paris, t. 333, Série I (2001), 81-86.
- [LR4] J.-L. Loday and M. Ronco, *Trialgebras and families of polytopes*, Preprint arXiv math.AT/0205043, to appear in Contemporary Maths. (2003).
- [MR] C. Malvenuto and Ch. Reutenauer, *Duality between quasi-symmetric functions and the Solomon descent algebra*, J. Algebra 177 (1995), no. 3, 967-982.
- [S] L. Solomon *A Mackey formula in the group ring of a Coxeter group*, J. Algebra 41 (1976), 255–264.